



Correction to: Casper Versus Precise Stent for the Treatment of Patients with Idiopathic Intracranial Hypertension

Nebiyat F. Belachew¹ · Severin Baschung² · William Almiri¹ · Ruben Encinas¹ · Johannes Kaesmacher^{1,3} · Tomas Dobrocky¹ · Christoph J. Schankin⁴ · Mathias Abegg⁵ · Eike I. Piechowiak¹ · Andreas Raabe⁶ · Jan Gralla¹ · Pasquale Mordasini¹

© The Author(s) 2021

Correction to:

Clin Neuroradiol 2021

<https://doi.org/10.1007/s00062-021-01024-2>

After the article was published online, the abstract was revised again. The correct abstract is given below. The original article has been corrected.

Abstract

Purpose We hypothesized that due to its specific characteristics, the CasperTM RX carotid stent (CP) might be particularly suitable for venous sinus stenting (VSS) in patients with idiopathic intracranial hypertension (IIH). To test this theory, we compared it to the commonly used Precise Pro RXTM stent (PP).

Methods A total of 15 patients with IIH (median age 28.7 years) were reviewed retrospectively. Technical aspects as well as peri- and postinterventional complication rates were examined in patients treated with CP ($n=10$) and the PP ($n=5$). Improvements in cerebrospinal fluid opening pressure (CSF OP), transstenotic pressure gradient (TSPG) and clinical symptoms were also assessed.

Results Stent delivery was easier and more successful with the CP than the PP (difficult/failed stent delivery 0.0% versus 57.1%) and consequently achieved with less attempts (≥ 2 : 0.0% versus 40.0%). No severe peri- or postinterventional complications or instances of in-stent thrombosis and/or stenosis were observed during follow-up. Improvement of CSF OP and TSPG immediately after VSS as well as at 6-month follow-up was comparable between the CP and PP group. Both groups showed substantial and similar decreases in intensity and frequency of headache. Almost all patients with other IIH-related symptoms showed either improvement or complete resolution of those symptoms after VSS. All patients who were available for interview ($n=12/15$) reported a substantial improvement in quality of life.

Conclusion VSS using the CP seems to be safe and effective. The CP may reduce the risk of difficult or failed stent delivery in patients with challenging intracranial venous anatomy.

Funding Open Access funding provided by University of Bern.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included

The online version of the original article can be found under <https://doi.org/10.1007/s00062-021-01024-2>.

✉ Nebiyat F. Belachew
nebiyatfilate.belachew@insel.ch

¹ Department of Diagnostic and Interventional Neuroradiology, Inselspital, Bern University Hospital, and University of Bern, Freiburgstraße 18, 3010 Bern, Switzerland

² Faculty of Medicine, University of Bern, Bern, Switzerland

³ Department of Diagnostic, Interventional and Pediatric Radiology, Inselspital, Bern University Hospital, and University of Bern, Bern, Switzerland

⁴ Department of Neurology, Inselspital, Bern University Hospital, and University of Bern, Bern, Switzerland

⁵ Department of Ophthalmology, Inselspital, Bern University Hospital, and University of Bern, Bern, Switzerland

⁶ Department of Neurosurgery, Inselspital, Bern University Hospital, and University of Bern, Bern, Switzerland

in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.